

ALGEBRA I PRELIM, DECEMBER 2013
(SEE THE NEXT PAGE)

(Analysis and Topology prelims follow)

1a) If $f : G \rightarrow G'$ is a group homomorphism and H is a subgroup of G , is it always the case that $f^{-1}(f(H)) = H$ (where $f^{-1}(f(H))$ denotes the inverse image of $f(H)$ in G or equivalently the set of all elements in G that map into $f(H)$)? Prove it or give a counterexample.

b) If the answer to part a is no, then give a complete description of the subgroup of G that equals $f^{-1}(f(H))$ and apply this description to your counterexample. If the answer to part a is yes, then you can skip this part of the question.

2) Let G^c denote the commutator subgroup of a group G .

a) Let N be any normal subgroup of G . Prove that G/N is abelian if and only if $G^c \subseteq N$.

b) Show if H is any subgroup of G such that $G^c \subseteq H$, then H is a normal subgroup of G .

c) Recall that a group G is called solvable if there exists a normal tower of subgroups starting with G and ending with $\{e\}$ for which all the factor groups are abelian. Now define the following subgroups of G , $G^{(0)} = G$, $G^{(1)} = G^c =$ commutator subgroup of G , $G^{(2)} = (G^{(1)})^c =$ commutator subgroup of $G^{(1)}$, $G^{(3)} = (G^{(2)})^c$, etc. Show that G is solvable if and only if $G^{(s)} = \{e\}$ for some positive integer s .

3a) Suppose that $K \trianglelefteq H \trianglelefteq G$ and K is a Sylow p -subgroup of H for some prime p , prove that $K \trianglelefteq G$.

b) Suppose G is a finite group, P is a Sylow p -subgroup, and H a (not necessarily Sylow) p -subgroup of G . If $H \subseteq N_G(P)$, prove that $H \subseteq P$.

4) Let A be an integral domain and M an A module. An element $x \in M$ is called a torsion element of M if the annihilator of x , $\text{Ann}(x)$, does not equal $\{0\}$.

(a) Show that the subset $T(M)$ of torsion elements of M forms a submodule of M . Would this still be the case, if A were not an integral domain? Prove it or give a counterexample.

(b) If $f : M \rightarrow N$ is a homomorphism of A -modules, show that $f(T(M)) \subseteq T(N)$.

(c) If $0 \rightarrow M' \xrightarrow{f} M \xrightarrow{g} M''$ is an exact sequence of A -modules, show that $0 \rightarrow T(M') \xrightarrow{f'} T(M) \xrightarrow{g'} T(M'')$ is also exact where f' and g' are the restrictions of f and g to $T(M')$ and $T(M)$, respectively.

(d) Would the result in (c) still be true if we added a 0 to the right-hand side of the above two sequences? Prove it or give a counterexample.

5) For each of the following, give an example if possible, and if not possible, **briefly** explain why not. All rings in this problem are assumed to be commutative and all of your examples should involve only commutative rings. *For purposes of the prelims, you must get at least 5 of these correct in order to have the problem count as being correct.*

(a) A commutative ring A and a multiplicatively closed subset S which does not contain 0, such that $S^{-1}(A)$ is not a local ring.

(b) A commutative ring A and a multiplicatively closed subset S which does not contain 0, such that all prime ideals of A intersect S ,

(c) A subring A of a Noetherian ring B that is not a Noetherian ring.

(d) A ring A and a submodule of a finitely generated A -module that is not finitely generated.

(e) A non-Noetherian ring A that is a Noetherian \mathbb{Z} -module.

(f) A Noetherian ring A that is a non Noetherian \mathbb{Z} -module.

Analysis Preliminary Exam, Fall 2013

Problem #1: Let $f : [0;1] \rightarrow \mathbb{R}$.

i) Define what it means for f to be Lipschitz. Also define what it means for f to be **absolutely** continuous.

ii) Prove that if f is Lipschitz, then it is absolutely continuous.

iii) Prove that $f(x) = \sqrt{x}$ is absolutely continuous, but not Lipschitz on $[0;1]$.

Problem #2: Define what it means for a function $f : [a;b] \rightarrow \mathbb{R}$ to be of bounded variation. Prove that a function of bounded variation is a difference of two monotone functions.

Problem #3: Recall that $E \subset \mathbb{R}^d$ is Lebesgue measurable if for every $\epsilon > 0$ there exists an open set O containing E such that $m(O \setminus E) < \epsilon$, where m is the outer measure. Prove that $E \subset \mathbb{R}^d$ is Lebesgue measurable if and only if for every $A \subset \mathbb{R}^d$,

$$m(A) = m(A \setminus E) + m(A \cap E):$$

Problem #4: Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be a continuous function on $[a;b]$ and twice continuously differentiable on $(a;b)$.

i) Suppose that $f''(x) > 0$ on $(a;b)$. Prove that

$$f((1-t)a + tb) \leq (1-t)f(a) + tf(b):$$

ii) Use i) to show that if $x, y \geq 0$ and $\frac{1}{p} + \frac{1}{q} = 1$, $p > 1$, then

$$xy \leq \frac{x^p}{p} + \frac{y^q}{q}:$$

iii) Use ii) to show that if f, g are absolutely integrable on \mathbb{R} , then

$$\int_{\mathbb{R}} f(x)g(x)dx \leq \left(\int_{\mathbb{R}} |f(x)|^p dx \right)^{\frac{1}{p}} \left(\int_{\mathbb{R}} |g(x)|^q dx \right)^{\frac{1}{q}}:$$

Problem #5: Let $f : \mathbb{R}^d \rightarrow \mathbb{C}$ be an absolutely integrable function. Prove that

$$\lim_{m \rightarrow \infty} \int_{\mathbb{R}^d} \frac{1}{1 + \frac{|x|^2}{m^2}} f(x) dx = \int_{\mathbb{R}^d} f(x) dx$$

I. Let A be a set. Let $\{X_j\}_j$ be an indexed family of spaces, and let $\{f_j: A \rightarrow X_j\}_j$ be an indexed family of functions $f_j: A \rightarrow X_j$.

(a) Show that there is a unique smallest (coarsest) topology T on A such that each f_j is continuous.

(b) Let $S = \{f_j^{-1}(U_j) \mid U_j \text{ is open in } X_j\}$ and $T = \langle S \rangle$. Show that S is a subbasis for the topology T on A .

(c) Show that a map $g: Y \rightarrow A$ is continuous relative to T if and only if each composition $f_j \circ g$ is continuous.

(d) Let $f: A \rightarrow \prod_j X_j$ be defined by

$$f(a) = (f_j(a))_j$$

and Z be the subspace $f(A)$ of the product space $\prod_j X_j$. Show that the image under f of each element of T is an open subset of Z .

II. Let X be a connected topological space. Let $x \in X$ be such that $X \setminus \{x\}$ is disconnected with separation $X \setminus \{x\} = A \cup B$. Prove $A \cup \{x\}$ is connected.

III. A topological space X is *countably compact* if every countable open covering of X contains a finite subcollection that covers X . Assume X is a Hausdorff space. Prove that the countably compact condition is equivalent to the (Bolzano-Weierstrass) condition that every infinite set in X has a limit point.

IV. Recall that a Baire space is characterized by the condition that any countable union of closed sets, each with empty interior, has empty interior. Prove that an open subspace of a Baire space is a Baire space.

V. Let $\{A_\alpha\}_\alpha$ be a locally finite collection of subsets of a topological space X . This condition means that each point of X lies in an open set which has a non-empty intersection with at most finitely many of the A_α . Prove

$$\overline{\bigcup_\alpha A_\alpha} = \bigcup_\alpha \overline{A_\alpha};$$