References

[Ber] B. C. Berndt, \Ramanujan's Notebooks I", Springer-Verlag, (1985).

[Br] R. Brown, \The Natural Density of Some Sets of Square-free Numbers."

[Ti] E. C. Titchmarsh, \The Theory of the Riemann Zeta Function." Second edition, published by Oxford University Press in 1986.

A Code

```
import math

def main():

    # Parameters and counts
    primes = [2, 3]       # List primes to filter out here
    N = 1000000             # Select number to count until
    square_free = 0       # Count for square frees
    sf_without_p = 0      # Count for square frees without the specified primes

    # Running the loop
    for i in range(N):

        # Variables for the loop
        square_free_add = 1                 # Assume i is square free
        without_add = 1                     # Assume i is indivisible by specified primes
        previous_prime = 1                  # Variable to keep track of previous prime
        cri = math.ceil(i ** (1 / 3))     # Highest factor we need to check is i^{1/3}

        while i > cri:
            next_prime = prime_factor(i)
            if next_prime == previous_prime:
                square_free_add = 0
                without_add = 0
                break
            if next_prime in primes:
                without_add = 0
            previous_prime = next_prime
            i /= previous_prime

        # Checking if remainder is equal to previous prime
        if i in primes:
            without_add = 0
        if i == previous_prime:
            square_free_add = 0
            without_add = 0

    # Print out results
    print(square_free)
    print(sf_without_p)

def prime_factor(n):
    if n in [1, 2, 3]:
        return n
    for i in range(2, math.ceil(math.sqrt(n)) + 1):
        if n % i == 0:
            return i
    return n
```