

Kapitza
Fall 2017

NOISY QUANTUM
MEASUREMENT

Ben Saltzman

Q.I.T. TERM PAPER

Professor Sarada Rajeev
December 10th 2017

1 Introduction

Not only does God play dice but... he sometimes throws them where they can't be seen.

Stephen Hawking

Despite Albert Einstein's insistence that the world is deterministic, quantum mechanics introduces a probabilistic world view. Measuring a quantum becomes a delicate task: measurements inherently affect the system in question. In this paper, based on a lecture given to fellow Kapitza members on December 3rd, 2017, I discuss the formulation of measurement in a noisy quantum system.

2 A New Description of Measurement

Measurement is commonly described by defining a set of projection operators that are complete, i.e.

\times

\sum_j

$P_j = |j\rangle\langle j|$

and the state of the system after measurement is

$$\frac{(I_S \otimes \text{tr}_{j^0} \rho_{j^0}^j)(U_{SP}^y \rho_{j^0}^j I_S \otimes j^0 i_P)}{\rho_j(j)} \quad (3)$$

Note that $I_S \otimes \text{tr}_{j^0} \rho_{j^0}^j$ is the measurement operator on the full system, since only the probe is measured.

Since U_{SP} is unitary, we have $U_{SP}^y U_{SP} = I_{SP} = I_S \otimes I_P$. Consider the case $k = 0$. For clarity, define $M^j = M_S^{j,0}$.

$$\begin{aligned} U_{SP}^y U_{SP} &\stackrel{k=0}{=} \prod_{j^0} M^{j^0,y} \otimes j^0 i_{h^0} j_P \prod_j M^j \otimes j j i_{h^0} j_P \\ &= \prod_{j^0} \prod_j M^{j^0,y} M^j \otimes j^0 i_{h^0} j j i_{h^0} j_P \\ &= \prod_j M^{j,y} M^j \otimes j i_{h^0} j_P \end{aligned} \quad (4)$$

But $U_{SP}^y U_{SP} \stackrel{k=0}{=} I_S \otimes j^0 i_{h^0} j_P$. Using (4), we conclude that

$$I_S = \prod_j M^{j,y} M^j \quad (5)$$

Using the definition of U_{SP} , we can simplify (2) and (3). Substituting (1) into (2):

$$\begin{aligned} \prod_j \rho_j(j) &= \prod_{j^0} \text{tr}_{j^0} \rho_{j^0}^j \prod_{j^0,k} M_S^{j^0,k,y} \otimes j^0 i_{h^0} j_P (I_S \otimes j j i_{h^0} j_P) \\ &\quad \times \prod_{j^0,k} M_S^{j^0,k} \otimes j j^0 i_{h^0} j_P \otimes j i_S \otimes j^0 i_P \\ &= \prod_{j^0,k} \text{tr}_{j^0} \rho_{j^0}^j M_S^{j^0,k,y} \otimes j^0 i_{h^0} j_P (I_S \otimes j j i_{h^0} j_P) \\ &\quad \times \prod_{j^0,k} M_S^{j^0,k} \otimes j i_S \otimes j j^0 i_{h^0} j_P \\ &= \prod_{j^0,k} \text{tr}_{j^0} \rho_{j^0}^j M_S^{j^0,y} I_S \otimes j j_P \prod_{j^0} M^{j^0} \otimes j i_S \otimes j j^0 i_P \\ &= \prod_j \text{tr}_{j^0} \rho_{j^0}^j M^{j,y} M^j \otimes j i \end{aligned}$$

and therefore

$$p_J(j) = \text{tr} [M_j^\dagger M_j \rho] \quad (6)$$

Similarly, we can simplify the post-measurement state to

$$\frac{M_j \rho_S \otimes \rho_P M_j^\dagger}{p_J(j)} \quad (7)$$

The state of S can be read off from (7) easily, since S and P are in a pure product state. Measurement can therefore be described as a set of measurement operators $\{M_j^S\}$, instead of $\{M_j\}$, that satisfy (5). See Appendix A for discussion on the application of this process to ensembles.

When transferring classical data over a quantum channel, the receiver doesn't need the post-measurement state to process the information in a quantum fashion. The relevant probability is the probability of error. For any such situation where the probability of an outcome matters and the post-measurement state does not, we can describe a positive operator-valued measure (POVM) with a set of operators $\{M_j^y\} = \{M_j^y M_j\}$ that are non-negative and complete. Clearly, projection is a type of POVM. The probability of success of the POVM is

$$\sum_{x \in X} p_X(x) \text{Tr}[\rho_X(x)]$$

where $\rho_X(x)$ is the density matrix for state $j = x$.

3 Composite Systems

Suppose we have two independent ensembles, $\rho_A = \sum_x p_X(x) |j_x\rangle\langle j_x|$ and $\rho_B = \sum_y p_Y(y) |j_y\rangle\langle j_y|$. The density matrix for the joint state $|j_x\rangle |j_y\rangle$ is

$$\begin{aligned} E_{X;Y}[(|j_x\rangle\langle j_x| \otimes |j_y\rangle\langle j_y|)(\rho_A \otimes \rho_B)] &= E_{X;Y}[|j_x\rangle\langle j_x| \otimes |j_y\rangle\langle j_y|] \\ &= \sum_x p_X(x) \sum_y p_Y(y) |j_x\rangle\langle j_x| \otimes |j_y\rangle\langle j_y| \\ &= \sum_x p_X(x) |j_x\rangle\langle j_x| \otimes \sum_y p_Y(y) |j_y\rangle\langle j_y| \\ &= \rho_A \otimes \rho_B \end{aligned} \quad (8)$$

where ρ_A and ρ_B are the density matrices for A and B respectively.

Now suppose we have a joint ensemble in which systems A and B are correlated classically. We'd like a formulation to express this ensemble similarly to the independent situation above. To do this, we introduce a new random variable Z that X and Y are conditioned on. The two ensembles

are $\rho_A = \int p_{X|Z}(x|z) |j_{x,z}\rangle\langle j_{x,z}|$ (density matrix ρ_A) and $\rho_B = \int p_{Y|Z}(y|z) |j_{y,z}\rangle\langle j_{y,z}|$ (density matrix ρ_B), and $X|Z$ and $Y|Z$ are independent. Using the same procedure as we did with (8), we obtain the density matrix of the total state:

$$\begin{aligned} E_{X,Y,Z}[(|j_{x,z}\rangle\langle j_{x,z}| \otimes |j_{y,z}\rangle\langle j_{y,z}|)(h_{x,z} \otimes h_{y,z})] \\ = \int_{x,y,z} p_Z(z) p_{X|Z}(x|z) p_{Y|Z}(y|z) |j_{x,z}\rangle\langle j_{x,z}| \otimes |j_{y,z}\rangle\langle j_{y,z}| \end{aligned} \quad (9)$$

Define a new random variable $W = X \wedge Y \wedge Z$. We can write the density matrix in (9) as

$$\int_W p_W(w) |j_w\rangle\langle j_w| \otimes |j_w\rangle\langle j_w| \quad (10)$$

So, we can write any state with the properties discussed in this paragraph as a product of pure states. This type of state is termed *separable*, and contains no entanglement. In other words, a separable state can always be prepared classically. See Appendix B for an application involving separable states.

4 Local Density Operators

Suppose systems A and B are in an entangled Bell state $|j\rangle_{AB}$. Take a POVM $\{J_j\}$ on A . The measurement operators for the system are $J_A^j \otimes I_B$. The probability of outcome j is

$$\begin{aligned} p_J(j) &= \langle j | J_A^j \otimes I_B | j \rangle_{AB} \\ &= \frac{1}{2} \sum_{k,l=0}^1 \langle k | \langle l | J_A^j | k \rangle | l \rangle_{AB} \\ &= \frac{1}{2} \sum_{k,l=0}^1 \langle k | J_A^j | l \rangle \langle l | J_A^j | k \rangle \\ &= \text{Tr}_A \left(J_A^j \frac{1}{2} \sum_{k=0}^1 |k\rangle\langle k| \right) \\ &= \text{Tr}_A \left(J_A^j \rho_A \right) \end{aligned} \quad (11)$$

where the "local density operator" for A is the maximally mixed state $\rho_A = \frac{1}{2} \sum_{k=0}^1 |k\rangle\langle k|$. This process goes similarly for B . Thus, the following global state gives the same predictions in local measurements as $|j\rangle_{AB}$:

$$\rho_A \otimes \rho_B$$

We'd like to define what we mean by a local density operator in order to describe the results of local measurements. To do this, we need to define the *partial trace* operation.

Suppose $\{f_j k_i\}_A$ and $\{f_l i_B\}_B$ are orthonormal bases for the Hilbert spaces of A and B . Then $\{f_j k_i\}_A \{f_l i_B\}_B$ is an orthonormal basis for the product of the Hilbert spaces. For density operator ρ_{AB} , the probability of outcome j is

$$\begin{aligned}
 p_J(j) &= \text{Tr} \left(\sum_{k,i} |f_j k_i\rangle \langle f_j k_i| \rho_{AB} \right) \\
 &= \sum_{k,i} \langle f_j k_i | \rho_{AB} | f_j k_i \rangle \\
 &= \sum_{k,i} \langle f_j | \langle k_i | \rho_{AB} | k_i \rangle | f_j \rangle \\
 &= \sum_{k,i} \langle f_j | \left(\sum_{l,m} |f_l i_B\rangle \langle f_l i_B| \rho_{AB} \right) | k_i \rangle | f_j \rangle \\
 &= \sum_{k,i} \langle f_j | \left(\sum_{l,m} |f_l i_B\rangle \langle f_l i_B| \rho_{AB} \right) | k_i \rangle | f_j \rangle \\
 &= \text{Tr}_B \left(\sum_{k,i} |f_j k_i\rangle \langle f_j k_i| \rho_{AB} \right) \\
 &= \text{Tr}_B \left(\sum_{k,i} |f_j k_i\rangle \langle f_j k_i| \rho_{AB} \right) \tag{12}
 \end{aligned}$$

So, we define the partial trace for B as

$$\text{Tr}_B[\rho_{AB}] = \sum_{l,i} |f_l i_B\rangle \langle f_l i_B| \rho_{AB} | f_l i_B\rangle \tag{13}$$

and the local density operator for A as

$$\rho_A = \text{Tr}_B[\rho_{AB}] \tag{14}$$

Therefore, (12) becomes

$$p_J(j) = \text{Tr}_A \left(\rho_A |f_j\rangle \langle f_j| \right) \tag{15}$$

Alice can predict the outcome of local measurements with (15).

5 Classical-Quantum Ensembles

Suppose Alice prepares a quantum system with density matrix ρ_A and probability distribution $p_X(x)$. She passes this ensemble to Bob, who must learn about it. There is a loss of information in X after preparation which is minimized if the state is pure. $\rho_A = \sum_x p_X(x) |x\rangle \langle x|$ for an orthonormal basis $\{|x\rangle\}_{x \in X}$. For a mixed state, $\rho_A = \sum_x p_X(x) |x\rangle \langle x|$ is more difficult to extract information from.

One solution is for Alice to prepare a *classical-quantum ensemble*:

$$\{p_X(x); |j\rangle\langle j|_X \otimes \rho_A^x\}_{x \in X}$$

This ensemble is so-called because system X is classical, while system A is quantum. The density operator for the entire system is

$$\rho_{XA} = \sum_x p_X(x) |j\rangle\langle j|_X \otimes \rho_A^x \quad (16)$$

Suppose Bob makes a measurement of the system with $\{I_X \otimes J_A^j\}$. This is akin to Bob measuring an isolated system A with $\{J_A^j\}$. Why?

$$\begin{aligned} \text{Tr}_{XA}(I_X \otimes J_A^j) &= \text{Tr} \left(\sum_x p_X(x) |j\rangle\langle j|_X \otimes \rho_A^x (I_X \otimes J_A^j) \right) \\ &= \text{Tr} \sum_x p_X(x) (|j\rangle\langle j|_X \otimes \rho_A^x) (I_X \otimes J_A^j) \\ &= \sum_x \text{Tr}[|j\rangle\langle j|_X \otimes \rho_A^x] \text{Tr} p_X(x) J_A^j \\ &= \sum_x \text{Tr} p_X(x) J_A^j \\ &= \text{Tr} J_A^j \end{aligned}$$

So Bob can extract information about A from the whole system with a local measurement on A .

6 Conclusion

At the heart of quantum mechanics is a rule that sometimes governs politicians or CEOs - as long as no one is watching, anything goes.

Lawrence M. Krauss

Measurement of quantum systems is tricky, and matters only get more complicated when you consider composite quantum systems. With judicious choices of measurement operators and careful preparation of a system or composite system, one can make quantum measurement seem more akin to the classical case.

Appendix A: Redefining Measurement for Ensembles

Suppose we have an ensemble

$$\rho_X(x); j \in \mathcal{X}$$

with density operator

$$\rho_X = \sum_{x \in \mathcal{X}} \rho_X(x) |x\rangle\langle x|$$

Using the same procedure as outlined in section 2:

$$\begin{aligned} \rho_J(j) &= \sum_{x \in \mathcal{X}} \rho_X(x) \langle j| \langle x|_S \rho_{j^0 j^1} \sum_{j^0, k} M_S^{j^0, ky} |j^0\rangle\langle j^1|_P (I_S \otimes |j\rangle\langle j|_P) \\ &= \sum_{x \in \mathcal{X}} \rho_X(x) \sum_{j^0, k} \langle j| \langle x|_S M_S^{j^0, ky} \langle j^0| \langle k|_P \rho_{j^1 j^0} (I_S \otimes |j\rangle\langle j|_P) \\ &= \sum_{x \in \mathcal{X}} \rho_X(x) \sum_{j^0, k} \langle j| \langle x|_S M_S^{j^0, ky} I_S \langle j^0| \langle k|_P \rho_{j^1 j^0} \\ &= \sum_{x \in \mathcal{X}} \rho_X(x) \sum_{j^0} \langle j| \langle x|_S M_S^{j^0, ky} \rho_{j^1 j^0} \\ &= \sum_{x \in \mathcal{X}} \rho_X(x) \langle j| \langle x|_S M_S^{j^0, ky} M_S^{j^0, ky} \rho_{j^1 j^0} \\ &= \text{Tr} [M_S^{j^0, ky} M_S^{j^0, ky}] \end{aligned}$$

This is a reformulation of the Born Rule. The post-measurement state is

$$M_S^{j^0, ky} \rho_{j^1 j^0}$$

