At Rochester, the Depart
Electrical and Computer
Science Foundation (NSF)
and awards to come by.
https://www.pas.rochest
quote Albert Einstein, “Th

9:00 AM

Andrew Hotchkiss, Houghton University

9:15 AM

Levi Kennel, Houghton University

9:30 AM

Timothy Ockrin, Houghton University

9:00 AM

Waly M Z Karim, University of Rochester

9:15 AM

Julia Largett, University of Rochester

9:30 AM

2:00 PM

Andrew Hotchkiss and Mark Yuly, Houghton University

The Houghton University cyclotron accelerates ions inside of a 17 cm inner diameter evacuated aluminum chamber placed between the poles of a 1.2 T electromagnet. Two hollow "dee" electrodes, one grounded and one oscillating between a positive and negative potential, apply an electric force that with the magnetic field, accelerates the ions. Previously, weak magnetic focusing, which requires the magnetic field to decrease with larger radius, was used to force the ions back toward the central plane between dees. For large numbers of orbits, however, this made the ions get out of phase with the oscillating electric field. Strong focusing can fix this problem by creating a restoring force while keeping the cyclotron frequency the same. Mathematica and Radia, a 3D magnetic field modeling software package, were used to model the magnetic field of a strong-focusing sector pole tip, then calculate resulting ion orbits. In simulations, the original weak focusing pole tips allowed an ion orbit radius of 3.7 cm to be reached. Using newly designed strong focusing pole tips a radius of 4.9 cm corresponding to 165 keV for protons was achieved. The maximum radius theoretically achievable is 7.2 cm resulting in 360 keV of proton energy.

Levi Kennel and Mark Yuly, Houghton University

Because there are currently no published weak interaction parity violation experiments specifically for undergraduate laboratories, a simple parity violation experiment is being developed using circularly polarized gamma rays. A 60Co source will be placed on one side of an electromagnet, so that the circularly polarized gamma rays emitted opposite the beta particles will pass through the electromagnet core. A NaI detector detects the number of gamma rays that pass through the electromagnet, and a silicon detector detects beta particles opposite the gamma rays. The number of coincidence events will be measured when the electromagnet is polarized both parallel and antiparallel to the gamma rays – an asymmetry between the number of coincidence events for each orientation would show that parity is violated.

Timothy Ockrin and Katrina Koehler, Houghton University

Decay Energy Spectroscopy (DES) results in high energy resolution (1-5 keV @ 5 MeV) spectra of decay energies where the energy of each decay is measured as a single event as opposed to individual measurements of each decay particle. In order to accomplish this, the measured source is not external to the absorber, but embedded within it. DES can be used for nuclear safeguards, metrology, and medical isotope development, but measurements are affected by incomplete energy capture occurring when decay particles escape the absorber. In order to reduce escape likelihood, absorbers can be capped with a layer of gold. Geant4, a Monte-Carlo simulation software capable of handling energy transport between particles, is used to simulate DES measurements with varying th

Waly M Z Karim and Segev BenZvi, University of Rochester

The IceCube Neutrino Observatory is capable of detecting high-energy astrophysical neutrinos as well as bursts of MeV neutrinos from core-collapse supernovae (CCSNe). The IceCube-Gen2 will encompass nearly tenfold the volume of its predecessor, incorporating cutting-incP0 (iMT0 (wi)4 (l)D6 (h)]T6)10 (o)t seng M1 (rs)2 (itl-4 (Z)-1

Delaney Cummins, Segev BenZvi and Xinyi Chen, University of Rochester

Strong gravitational lensing systems are typically observed in imaging surveys via rings, multiple images, or other visual effects on the source galaxy. However, we expect to see 5 to 10 times more strong lensing systems with Einstein radii smaller than the resolution of most imaging surveys. These systems may be found using spectroscopy, and their detection would dramatically increase the number of known lensing systems, improving our chances to perform time-delay cosmography and other cosmologically significant studies. We train a vision transformer neural network (ViT) to find strong lensing systems in DESI spectra. The ViT classified lenses and non-lenses with >85% accuracy after training.

Kylie L. Goldade, Victoria S. Wynter, Jason P. Bier, Matthew J. Wright, Adelphi University

We will report our findings about doing physics outreach through Adelphi University. We do various events through Lab4Kids which is a program that allows us to travel to high schools to teach students, as well as them being able to visit Adelphi. In this program, we taught children (grades K - 12) in groups of up to 32 students following the NYS Curriculum and beyond, traveled to high schools, and brought students back to our nationally recognized STEM University. Because of the opportunities from this program, three of our student teachers were invited to teach for the 21st Century Community Learning Centers RamPathways Program at the West Hempstead School District. This program allowed us to teach STEM Enrichment during the summer following a common core aligned curriculum, designing, writing, and implementing lessons for subjects including optics, electrostatics, electromagnetism, and forces, as well as modifying assignments for students with disabilities and special needs. Victoria Wynter taught 12, 5th-grade students at George Washington Elementary School. Kylie Goldade taught 23, 6th-grade students at George Washington Elementary. Jason Bier taught 15, 7th, and 8th grade students at West Hempstead Secondary School.

Mimi Harrison and Jeyhan Kartaltepe, Rochester Institute of Technology

Massive galaxies exhibit a bimodal distribution in the Universe, consisting of active star-forming and quiescent populations persisting through high redshifts. In a cosmologically young Universe only a few billion years old, it is a mystery how massive galaxies could have become quenched of star formation in such significant numbers so quickly. This phenomenon is not predicted nor explained by theoretical models of galaxy evolution and hierarchical structure formation models- suggesting that galaxy evolution in the early Universe may differ dramatically from what we observe in the current Universe. With access to revolutionary observational equipment, exploring these rare galaxies from the early Universe is possible with larger sample sizes than previously possible. We present a sample of quiescent galaxy candidates selected from the COSMOS-Web catalogs by using rest-frame color-color diagrams with F115W, F150W, F277W, F444W, and F770W filters to identify these galaxies and distinguish them from active star-forming ones. Spectral Energy Distribution (SED) models from photometric COSMOS-Web data can test whether these candidates fit typical quiescent galaxies' SED profiles. Future steps to utilize spectroscopic quiescent candidate galaxy data from the Keck Observatory will allow for further study and test the efficacy of color-color techniques at higher redshifts.

Declan Walden, Debamitra Chakraborty, Jing Cheng, Antanas Straksys, Arunas Stirke, Wanessa CMA Melo, Arturas Jukna, Ivan V. Komissarov, and Roman Sobolewski, University of Rochester

Gelatin exhibits excellent biodegradability, biocompatibility, adhesion, and absorption properties. This polymer can undergo varying degrees of cross-linking with glutaraldehyde (GTA) and linked with water to form a hydrogel. The water diffusion coefficient serves as an important parameter governing the applicability of hydrogels. Various techniques can be employed to measure water diffusion in gelatin hydrogels, including quasi-elastic neutron scattering, nuclear magnetic resonance, side-by-side diffusion cells, fluorescence correlation spectroscopy, the refractive index method, among others. In this study, we introduce a novel method for determining water diffusion in gelatin hydrogels based on transient terahertz time-domain spectroscopy (THz-TDS). The diffusion coefficient is derived from the temporal evolution of experimentally measured water-concentration profiles using THZ-TDS. Assuming Fickian diffusion, the water diffusion coefficient is estimated to be 2.53 * 10^–6 cm2/s, falling within the range of reported values for water diffusion coefficients in gelatin literature.

Corey J. Young, Gable M. Wadsworth and Priya R. Banerjee, SUNY Buffalo

The phase separation of RNA and proteins is ubiquitous in biological systems, an important example being the formation of membrane-less ribonucleoprotein organelles such as the stress granules and the nucleolus in living cells. The majority of the past and current studies are focused on understanding the molecular rules of phase separation of proteins. RNAs can also undergo phase transitions in the absence of proteins. However, the physicochemical driving forces of protein-free, RNA-driven phase transitions remain unclear. We recently demonstrated RNAs undergo lower critical solution temperature (LCST)-type phase separation upon heating (Nature Chemistry, 2023; DOI: 10.1038/s41557-023-01353-4), which is driven by the desolvation of the phosphate backbone in the presence of Mg+2 ions and tuned by nucleobases. Here we uncover the potential for systematic perturbation of the RNA phase separation. Selecting a simple yet diverse set of parameters, such as divalent ions, molecular crowding, and sequence variation to tune RNA phase behavior, our work establishes novel heuristics to encode progra

Shaib Sharhan, Manhattan College

This research project, conducted at the European Organization for Nuclear Research (CERN) during the summer, aimed to advance the field of neutrino detection through innovative methodolog0.003 Tci.003 TT (he)13 ()TJ0 (s)6 (a)14 (e)]TJ0 Tc 0 Tw 15923 0 Td(-)Tj0.003 Tc -
nUNttttrc(r)4 c(r)4h(C)-4 (a)10rg(o)2e(tt)6,(g)1 (Ap)-4lay(r)4 ( )10a f TT uApCnh5 -9tt 6Ctthneottgl914tAp
U dergo NNneutrin1210 ( )]TJT* (E)3x(c)8p(de)3 (r)4 (i)4 (m)4 (e)13nt [c)18D[U)7 (N)1 (E)3 ())8 (a)4 (i)4 (
-5(o)21(1210 t(pe)3 (a)14m a)4 (t)10 (o)2(d o)12pt(he)9ia1314 ttaUNEn,

Allison Blum, Ka-Wah Wong, Rodrigo S. Nemmen, Jimmy Irwin and Dacheng Lin, SUNY Brockport

Most of the active galactic nuclei (AGNs) in the local universe are low-luminosity AGN (LLAGNs). They are generally associated with radiatively inefficient accretion flows (RIAFs). We present our study of the gamma-ray emitting radio galaxy NGC 315 hosting a LLAGN with a misaligned jet. Multiwavelength studies suggest that X-rays come from jets, however, it is also possible that the RIAF can contribute significantly to the X-rays. We present our study of the origin of the X-rays by using the NuSTAR X-ray Observatory. Using our X-ray data, we also provide constrains on the synchrotron self-Compton (SSC) model, which is believed to be the gamma-ray emitting mechanism for the LLAGN. We further study the origin of the X-ray emission using the Chandra X-ray Observatory, utilizing the high-resolution images created by the deconvolution method.

Sean Lannon, Zachary Robinson, Matt Sullivan, Karsten Beckmann and Nate Cady, SUNY Brockport

Niobium oxide is a strong candidate for implementation into neuromorphic computer technology. The purpose of this research is to develop synthesis and crystallization processes for making thin films of niobium oxide so that the material can be better understood and eventually incorporated into next-generation computers. More specifically, we studied the environmental conditions required for the material to undergo crystallization. We built a vacuum system with a regulated inflow of gas and a furnace so that we could control the temperature and pressure of the sample's environment. The samples were measured using microscopic image analysis and Raman spectroscopy. We found that the samples underwent the most crystallization at pressures around atmospheric pressure, and temperatures around 900 . Crystallization only occurred when using an ambient of nitrogen/hydrogen gas mixture at 95%/5%. Crystallization did not consistently occur in(t)o1 (n()18 (d ( )10 (u)-4 ((o)2 (c5 Tc 1(%

Avery Gilson and Dr. Ben Ovryn, New York Institute of Technology

Using a broadband electro-optic phase modulator with a laser feedback interferometer enables measurements with phase shifting and lock-in approaches. We have calibrated the response of a piezoelectric translator and measured the dynamic response of a 5-axis robotic arm. A field programmable gate array (FPGA) offers the benefit of rapid phase measurements as well as closed loop control. Phase shifting laser feedback interferometry readily produces kHz phase measurements with better than 5 nm precision. Phase unwrapping can be used to follow micron changes in the optical path length. Our calibration of a commercial piezoelectric translator is in close agreement with vendor specifications. Stepper motors on the robotic arm combined with kinematic calculations produce accurate, large scale sample displacements (centimeter scale) while the laser feedback interferometer adds nanometer precision.

Amii Matamoros, Arie Bodek, M.E. Christy and Zihao Lin, University of Rochester

We report on parameterizations of nuclear excitation longitudinal and transverse nuclear electromagnetic form factors measured in electron scattering on 27 Al nuclei. For the elastic and nuclear excitations with excitation energies below 8 MeV we use all published form factors. These parameterizations can be used to calculate radiative corrections for electron scattering cross sections on 27 Al nuclear targets.

Russell Berger, Alex Mavian, Nazifa Rumman, Pascal Bassène, Humberto Terrones and Moussa N'Gom, Rensselaer Polytechnic Institute

Two-dimensional transition metal dichalcogenides (TMDs) are exciting candidates for miniaturizing nonlinear optical systems due to their high second-order nonlinear susceptibility. However, strong second harmonic generation (SHG) responses are currently hindered by their nanometer-scale interaction length, compounded by other physical factors, such as grain boundaries and multilayer growth, which limit the spatial degrees of freedom for signal generation. We incorporate wavefront shaping (WFS) techniques to enhance the weak SHG response of few-layered WS2 samples. Star-shaped polycrystals are examined for their weak SHG signal at both grain boundaries and the center of the crystal. We observe a significant targeted enhancement of the second-order nonlinear response in selected regions by up to an order of magnitude and report an increased conversion efficiency in these regions. Our work demonstrates how structured light can amplify weak nonlinear interactions, allowing for all-optical tuning of the nonlinear optical response in TMDs and paving the way for applications in nonlinear microscopy, nanophotonics, and materials characterization.

Name	Status	Institution
Sumedh Anantha	Undergraduate Student	University of Rochester
Nur Ashsams	Undergraduate Student	University at Buffalo
Divin Bayubahe	Undergraduate Student	Roberts Wesleyan University
Avery Belanger	Undergraduate Student	Houghty[illegible] at

Faraz Khan	Undergraduate Student	New York Institute of Technology
Masroor Khonkhodzhaev	Undergraduate Student	CUNY Queens College
Will Kittelberger	Undergraduate Student	University of Rochester
Asher Knight	Undergraduate Student	Houghton University
Katrina Koehler	Faculty	Houghton University
Sean Lannon	Undergraduate Student	SUNY Brockport
Julia Largett	Undergraduate Student	University of Rochester
Chunsun Lei	Undergraduate Student	Houghton University
Annie Maloney	Undergraduate Student	University of Rochester
Andrew Martin	Undergraduate Student	Houghton University


Genesee River
Prince Athletic Complex
Fauver Stadium
Goergen Athletic Center
Lyman Squash and Racquet Center
Speegle-Wilbraham Aquatic Center
Gilbert
Spurrier
Sage
Crosby
Dandelion Square
Susan B. Anthony Bowl
Wilson Quadrangle
Douglass Leadership House
Burton
Wilson Commons
Douglass Commons
University Health Service
Drama House
LeChase
Rettner
Lechase Courtyard
Rush Rhees Library
Todd
Lattimore
Strong Auditorium
River Campus
Bausch & Lomb
Meliora
Dewey
Hoyt
Schlegel
Carol G. Simon
Harkness
Interfaith Chapel
Gleason
Gavett
Hopeman
Wallis
Wegmans
New York State Optics
Genesee Valley Park
Hajim School of Engineering Quadrangle
Wilmot
Hylan
Goergen
Hutchison
Mount Hope
Visitor Info
Pump Station
Trustee Rd.
Trustee Lot
Ice Arena
Central Utilities